

The GAMMA-400 gamma-ray telescope for precision gamma-ray emission investigations

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2016 J. Phys.: Conf. Ser. 675 032009

(<http://iopscience.iop.org/1742-6596/675/3/032009>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 46.39.49.101

This content was downloaded on 07/02/2016 at 08:11

Please note that [terms and conditions apply](#).

The GAMMA-400 gamma-ray telescope for precision gamma-ray emission investigations

N P Topchiev^{1,*}, A M Galper^{1,2}, V Bonvicini³, O Adriani⁴, R L Aptekar⁵, I V Arkhangel'skaja², A I Arkhangel'skiy², A V Bakaldin², L Bergstrom⁶, E Berti⁴, G Bigongiari⁷, S G Bobkov⁸, M Boezio³, E A Bogomolov⁵, L Bonechi⁷, M Bongi⁴, S Bottai⁴, G Castellini⁹, P W Cattaneo¹⁰, P Cumani^{3,16}, O D Dalkarov¹, G L Dedenko², C De Donato¹¹, V A Dogiel¹, N Finetti⁴, D Gascon¹², M S Gorbunov⁸, Yu V Gusakov¹, B I Hnatyk¹³, V V Kadilin², V A Kaplin², A A Kaplun², M D Kheyimits², V E Korepanov¹⁴, J Larsson¹⁵, A A Leonov^{1,2}, V A Loginov², F Longo³, P Maestro⁷, P S Marrocchesi⁷, M Martinez¹⁶, A L Men'shenin¹⁷, V V Mikhailov², E Mocchiutti³, A A Moiseev¹⁸, N Mori⁴, I V Moskalenko¹⁹, P Yu Naumov², P Papini⁴, J M Paredes¹², M Pearce¹⁵, P Picozza¹¹, A Rappoldi¹⁰, S Ricciarini⁹, M F Runtso², F Ryde¹⁵, O V Serdin⁸, R Sparvoli¹¹, P Spillantini⁴, Yu I Stozhkov¹, S I Suchkov¹, A A Taraskin², M Tavani²⁰, A Tiberio⁴, E M Tyurin², M V Ulanov⁵, A Vacchi³, E Vannuccini⁴, G I Vasilyev⁵, J E Ward¹⁶, Yu T Yurkin², N Zampa³, V N Zirakashvili²¹ and V G Zverev²

¹ Lebedev Physical Institute, RU-119991 Moscow, Russia

² National Research Nuclear University MEPhI (Moscow Engineering Physics Institute), Kashirskoe highway 31, Moscow, 115409, Russia

³ Istituto Nazionale di Fisica Nucleare, Sezione di Trieste, I-34149 Trieste, Italy

⁴ Istituto Nazionale di Fisica Nucleare, Sezione di Florence, I-50019 Sesto Fiorentino, Florence, Italy

⁵ Ioffe Physical Technical Institute, RU-194021 St. Petersburg, Russia

⁶ The Oskar Klein Centre, Department of Physics, Stockholm University, AlbaNova University Center, SE-106 91 Stockholm, Sweden

⁷ Department of Physical Sciences, Earth and Environment, University of Siena and Istituto Nazionale di Fisica Nucleare, Sezione di Pisa, Italy

⁸ Scientific Research Institute for System Analysis, RU-117218 Moscow, Russia

⁹ Istituto di Fisica Applicata Nello Carrara, I-50019 Sesto Fiorentino, Florence, Italy

¹⁰ Istituto Nazionale di Fisica Nucleare, Sezione di Pavia, I-27100 Pavia, Italy

¹¹ Istituto Nazionale di Fisica Nucleare, Sezione di Rome "Tor Vergata", I-00133 Rome, Italy

¹² Departament d'Astronomia i Meteorologia, Institut de Ciències del Cosmos, Universitat de Barcelona, Spain

¹³ Taras Shevchenko National University, Kyiv, 01601 Ukraine

¹⁴ Lviv Center of Institute of Space Research, Lviv, 79060 Ukraine

¹⁵ KTH Royal Institute of Technology, Department of Physics and the Oskar Klein Centre, AlbaNova University Center, SE-10691 Stockholm, Sweden

¹⁶ Institut de Física d'Altes Energies, Bellaterra, Spain

¹⁷ Research Institute for Electromechanics, RU-143502 Istra, Moscow region, Russia



¹⁸ NASA Goddard Space Flight Center and CRESST/University of Maryland, Greenbelt, Maryland 20771, USA

¹⁹ Hansen Experimental Physics Laboratory and Kavli Institute for Particle Astrophysics and Cosmology, Stanford University, Stanford, CA 94305, USA

²⁰ Istituto Nazionale di Astrofisica IASF and Physics Department of University of Rome Tor Vergata, I-00133 Roma, Italy

²¹ Pushkov Institute of Terrestrial Magnetism, Ionosphere, and Radiowave Propagation, Troitsk, Moscow region, Russia

E-mail: *tnp51@yandex.ru

Abstract. The GAMMA-400 gamma-ray telescope with excellent angular and energy resolutions is designed to search for signatures of dark matter in the fluxes of gamma-ray emission and electrons + positrons. Precision investigations of gamma-ray emission from Galactic Center, Crab, Vela, Cygnus, Geminga, and other regions will be performed, as well as diffuse gamma-ray emission, along with measurements of high-energy electron + positron and nuclei fluxes. Furthermore, it will study gamma-ray bursts and gamma-ray emission from the Sun during periods of solar activity. The GAMMA-400 energy range is expected to be from ~20 MeV up to TeV energies for gamma rays, up to 10 TeV for electrons + positrons, and up to 10^{15} eV for cosmic-ray nuclei. For 100-GeV gamma rays, the GAMMA-400 angular resolution is $\sim 0.01^\circ$ and energy resolution is $\sim 1\%$; the proton rejection factor is $\sim 5 \times 10^5$. GAMMA-400 will be installed onboard the Russian space observatory.

1. Introduction

The main goal for the GAMMA-400 mission is to perform a sensitive search for signatures of dark matter particles in high-energy gamma-ray emission. This task was set for the GAMMA-400 project by Nobel Laureate Academician V.L. Ginzburg in the end of 1980's [1, 2] and in his list of very important issues in modern cosmology at the beginning of XXI Century noted the issue of dark matter and its detection [3]. Within the framework of this project, which has now since expanded internationally, the design and construction of a future, complex, precision gamma-ray telescope is being carried out [4-7].

It should be noted, up to now, neither in indirect measurements of EGRET [8], currently operating Fermi-LAT [9], ground-based gamma-ray facilities [10] nor in direct measurements of various ground-based experiments including experiments at LHC, any direct evidence of hypothetical dark matter particles were not found.

GAMMA-400 is designed to search for signatures of dark matter in the fluxes of gamma-ray emission and electrons + positrons. Precision investigations of gamma-ray emission from Galactic Center, Crab, Vela, Cygnus, Geminga, and other regions will be performed, as well as diffuse gamma-ray emission, along with measurements of high-energy electron + positron and nuclei fluxes. Furthermore, it will study gamma-ray bursts and gamma-ray emission from the Sun during periods of solar activity.

2. The GAMMA-400 gamma-ray telescope

The GAMMA-400 physical scheme is shown in figure 1. GAMMA-400 consists of plastic scintillation anticoincidence top and lateral detectors (AC_{top} and AC_{lat}), converter-tracker (C), plastic scintillation detectors (S1 and S2) for time-of-flight system (ToF), two-part calorimeter (CC1 and CC2), lateral detectors (LD), plastic scintillation detectors (S3 and S4), and neutron detector (ND).

The anticoincidence detectors surrounding the converter-tracker are used to distinguish gamma quanta from significantly larger number of charged particles (e.g., in the region of 10-100 GeV, the flux ratios for gamma rays to electrons and protons are $\sim 1:10^2:10^5$ [11]). The gammas recognizing

provides by the counting and triggers signals formation system due special algorithms of backslash rejection [12].

All scintillation detectors consist from two independent 1-cm layers. The time-of-flight system, where detectors S1 and S2 are separated by approximately 500 mm, determines the top-down direction of arriving particles. The additional scintillation detectors S3 and S4 improve hadron and electromagnetic shower separation.

The converter-tracker consists of 13 layers of double (x, y) silicon strip coordinate detectors (pitch of 0.08 mm). The first three and final two layers have no tungsten while the middle eight layers are interleaved with tungsten conversion foils. Using the first three layers without tungsten allows us to measure gamma rays down to approximately 20 MeV. The total converter-tracker thickness is about $1 X_0$ (where X_0 is the radiation length). The converter-tracker information is used to precisely determine the conversion point and the direction of each incident particle.




Figure 1. The GAMMA-400 physical scheme.

The two-part calorimeter measures particle energy. The imaging calorimeter CC1 consists of 2 layers of double (x, y) silicon strip coordinate detectors (pitch of 0.08 mm) interleaved with planes from CsI(Tl) crystals, and the electromagnetic calorimeter CC2 consists of CsI(Tl) cubic crystals with dimensions of $36 \text{ mm} \times 36 \text{ mm} \times 36 \text{ mm}$. The thickness of CC1 and CC2 is $2 X_0$ and $23 X_0$, respectively. The total calorimeter thickness is $25 X_0$ or $1.2 \lambda_0$ (where λ_0 is nuclear interaction length) when detecting vertical incident particles and $54 X_0$ or $2.5 \lambda_0$ when detecting laterally incident particles. Using a deep calorimeter allows us to extend the energy range up to several TeV for gamma rays, 10 TeV for electrons, and to reach an energy resolution of approximately 1% above 100 GeV.

Neutron detector is used to separate hadron and electromagnetic showers.

The GAMMA-400 initial steps to modeling of principal possibilities to detect particles from vertical and lateral directions are shown in figure 1 for several fields of view: (1) for gamma rays and electrons + positrons in the main aperture, (2) for gamma-rays and electrons + positrons in the additional one; and for gamma-rays, electrons + positrons, and nuclei in the lateral aperture (3). The results shows the best angular resolution of $\sim 0.01^\circ$ within $\pm 30^\circ$ for vertical particles ($E_\gamma = 100 \text{ GeV}$) in

the main aperture and $\sim 0.2^\circ$ for gammas with $E_\gamma = 100$ GeV in the additional one [13] and the possibility to extend energy range up to several TeV for gamma rays, 10 TeV for electrons in the lateral aperture [12].

3. The GAMMA-400 performance and comparison with Fermi-LAT and ground-based facilities

The GAMMA-400 performance has been presented earlier in [4-7]. At present, using additional first three double layers (without tungsten) allows us to achieve the best angular resolution in the energy range from ~ 20 MeV up to TeV in spite of some improved the Fermi-LAT performance with Pass 8 [14].

Table 1. Comparison of the Fermi-LAT and GAMMA-400 parameters.

	Fermi-LAT	GAMMA-400
Orbit	Circular, 565 km	Highly elliptical, 500-300000 km (without the Earth's occultation)
Operation mode	Sky-survey (3 hours)	Point observation (up to 100 days)
Source exposition	1/7	1
Energy range	20 MeV - 300 GeV (γ , e)	~ 20 MeV - 1 TeV (γ) 1 GeV - 10 TeV (e)
Effective area ($E_\gamma > 1$ GeV)	~ 6500 cm ² (total) ~ 4000 cm ² (front)	~ 4000 cm ²
Coordinate detectors - readout	Si strips (pitch 0.23 mm) digital	Si strips (pitch 0.08 mm) analog
Angular resolution	$\sim 4^\circ$ ($E_\gamma = 100$ MeV) $\sim 0.2^\circ$ ($E_\gamma = 10$ GeV) $\sim 0.1^\circ$ ($E_\gamma > 100$ GeV)	$\sim 2^\circ$ ($E_\gamma = 100$ MeV) $\sim 0.1^\circ$ ($E_\gamma = 10$ GeV) $\sim 0.01^\circ$ ($E_\gamma > 100$ GeV)
Calorimeter - thickness	CsI(Tl) $\sim 8.5 X_0$	CsI(Tl) + Si $\sim 25 X_0$
Energy resolution	$\sim 10\%$ ($E_\gamma = 10$ GeV) $\sim 10\%$ ($E_\gamma > 100$ GeV)	$\sim 3\%$ ($E_\gamma = 10$ GeV) $\sim 1\%$ ($E_\gamma > 100$ GeV)
Proton rejection factor	$\sim 10^3$	$\sim 5 \times 10^5$
Mass, kg	2800	4100
Telemetry downlink volume	15 Gbytes/day	100 Gbytes/day

Table 1 presents a comparison of the Fermi-LAT and GAMMA-400 parameters. GAMMA-400 has numerous advantages:

- highly elliptical orbit (without the Earth's occultation and away from the radiation belts) allows us to observe with the full aperture of $\pm 60^\circ$ different gamma-ray sources continuously over a long period of time with the exposition greater by a factor of 7 than for Fermi-LAT operating in the sky-survey mode;
- thanks to a smaller pitch (by a factor of 3) and analog readout in the coordinate silicon strip detectors, GAMMA-400 has an excellent angular resolution above ~ 20 MeV;
- due to the deep ($\sim 25 X_0$) calorimeter, GAMMA-400 has an excellent energy resolution and can more reliably to detect gamma rays up to several TeV and electrons + positrons up to 10 GeV for vertically incident events. When measuring lateral events (electrons + positrons and nuclei) it can detect diffuse gamma-ray emission and cosmic rays using only the CC2 calorimeter;
- owing to the better gamma-ray separation from cosmic rays (in contrast to Fermi-LAT, the presence of a special trigger with event timing, time-of-flight system, two-layer scintillation detectors), GAMMA-400 is significantly well equipped to separate gamma rays from the background of cosmic rays and backscattering events.

GAMMA-400 will have also the better angular and energy resolutions in the energy region 10-1000 GeV in comparison with current and future space- and ground-based instruments: H.E.S.S. [15], MAGIC [16], VERITAS [17], HAWC [18], CTA [19], DAMPE [20], and CALET [21] (figure 2, table 2) and it allows us to fill the gap between the space- and ground-based instruments.

Table 2. Comparison of parameters for the current, future space- and ground-based instruments.

	SPACE-BASED				GROUND-BASED			
	GAMMA-RAY INSTRUMENTS				GAMMA-RAY INSTRUMENTS			
	Fermi-LAT	DAMPE	CALET	GAMMA-400	H.E.S.S.	MAGIC	VERITAS	CTA
Particles	γ, e	e, nuclei, γ	e, nuclei, γ	$\gamma, e,$ nuclei	γ	γ	γ	γ
Operation period	2008-	2015	2015	~2023	2012-	2009-	2007-	~2020
Energy range, GeV	0.02-300	5-10000	10-10000	0.02-10000	> 30	> 50	> 100	> 20
Angular resolution ($E_\gamma > 100$ GeV)	0.1°	0.1°	0.1°	~0.01°	0.07°	0.07° ($E_\gamma = 300$ GeV)	0.1°	0.1° ($E_\gamma = 100$ GeV) 0.03° ($E_\gamma = 10$ TeV)
Energy resolution ($E_\gamma > 100$ GeV)	10%	1.5%	2%	~1%	15%	20% ($E_\gamma = 100$ GeV) 15% ($E_\gamma = 1$ TeV)	15%	20% ($E_\gamma = 100$ GeV) 5% ($E_\gamma = 10$ TeV)




Figure 2. Comparison of energy and angular resolutions for GAMMA-400, Fermi-LAT, H.E.S.S., HAWC, and CTA (figure 2 from [24] was used).

GAMMA-400 will study continuously over a long period of time different regions of Galactic plane, for example, Galactic Center, Crab, Vela, Cygnus, Geminga with the full aperture of $\pm 60^\circ$. In particular, using the gamma-ray fluxes obtained by Fermi-LAT [22, 23], we can expect that GAMMA-400 when

observing the Galactic Center with aperture of $\pm 45^\circ$ during 1 year will detect: 57400 photons for $E_\gamma > 10$ GeV; 5240 photons for $E_\gamma > 50$ GeV; 1280 photons for $E_\gamma > 100$ GeV; 535 photons for $E_\gamma > 200$ GeV.

4. GAMMA-400 gamma-ray observatory

In addition to the gamma-ray telescope, the GAMMA-400 scientific complex includes the KONUS-FG gamma-ray burst monitor and two star sensors for determining the gamma-ray telescope axis with an accuracy of $\sim 5''$, along with two magnetometers for measuring the magnetic field. The GAMMA-400 gamma-ray observatory will be installed onboard of the Navigator space platform, which is designed and manufactured by the Lavochkin Association.

Using the Navigator space platform gives the GAMMA-400 experiment a highly unique opportunity for the near future gamma- and cosmic-ray science, since it allows us to install a very large scientific payload (a mass of 4100 kg, power consumption of 2000 W, and a telemetry downlink of 100 GB/day, with lifetime more than 7 years), which will provide GAMMA-400 with the means to significantly contribute as the next generation instrument for gamma-ray astronomy and cosmic-ray physics.

The GAMMA-400 experiment will be initially launched into a highly elliptical orbit (with an apogee of 300,000 km and a perigee of 500 km, with an inclination of 51.4°), with 7 days orbital period. Under the action of gravitational disturbances of the Sun, Moon, and the Earth after ~ 6 months the orbit will transform to about an approximately circular one with a radius of $\sim 200,000$ km and will not suffer from the Earth's occultation and shielding by the radiation belts. A great advantage of such an orbit is the fact that the full sky coverage will always be available for gamma-ray astronomy, since the Earth will not cover a significant fraction of the sky, as is usually the case for low Earth orbit. Therefore, the GAMMA-400 source pointing strategy will hence be properly defined to maximize the physics outcome of the experiment. The launch of the GAMMA-400 space observatory is planned for the middle of the 2020s.

References

- [1] Dogiel V *et al.* 1988 *Space Sci. Rev.* **49** 215
- [2] Ginzburg V *et al.* 1995 *Adv. Space Res.* **15** 93
- [3] Ginzburg V 2004 *Phys. Usp.* **47** 1155
- [4] Galper A *et al.* 2013 *Adv. Space Res.* **51** 297
- [5] Galper A *et al.* 2013 *AIP Conf. Proc.* **1516** 288
- [6] Adriani O *et al.* 2014 *arXiv*:1412.4239
- [7] Topchiev N *et al.* 2015 *Bull. RAS. Physics* **79** 417
- [8] Wim de Boer 2005 *CERN COURIER* 29491
- [9] Ackermann M *et al.* 2015 *arXiv*:1506.00013
- [10] Doro M 2014 *arXiv*:1404.5017
- [11] Moiseev A A *et al.* 2007 *Astroparticle Phys.* **27** 339
- [12] Arkhangelskaja I V *et al.* 2015 *Physics Procedia* **74** 212
- [13] Leonov A A *et al.* 2015 *Physics Procedia* **74** 183
- [14] URL: http://www.slac.stanford.edu/exp/glast/groups/canda/lat_Performance.htm
- [15] Balzer A *et al.* 2013 *arXiv*:1311.3486
- [16] Mazin D *et al.* 2014 *arXiv*:1410.5073
- [17] Ong R E *et al.* 2014 *Adv. Space Res.* **53** 1483
- [18] Westerhoff S *et al.* 2014 *Adv. Space Res.* **53** 1492
- [19] CTA Consortium 2011 *Experimental Astronomy* **32** 93
- [20] Dong Yi-Fan *et al.* 2015 *arXiv*:1503.00415
- [21] Mori N 2013 *Nucl. Phys. B (Proc. Suppl.)* **239-240** 199
- [22] Ackermann M *et al.* 2013 *arXiv*:1306.6772
- [23] URL: <http://fermi.gsfc.nasa.gov/ssc/data/access/>
- [24] Funk S *et al.* 2013 *Astroparticle Phys.* **43** 348